

THE LAGRANGIAN OF q -POINCARÉ GRAVITY

Leonardo Castellani

Istituto Nazionale di Fisica Nucleare, Sezione di Torino
and
Dipartimento di Fisica Teorica
Via P. Giuria 1, 10125 Torino, Italy.

Abstract

The gauging of the q -Poincaré algebra of ref. [1] yields a non-commutative generalization of the Einstein-Cartan lagrangian.

We prove its invariance under local q -Lorentz rotations and, up to a total derivative, under q -diffeomorphisms. The variations of the fields are given by their q -Lie derivative, in analogy with the $q = 1$ case. The algebra of q -Lie derivatives is shown to close with field dependent structure functions.

The equations of motion are found, generalizing the Einstein equations and the zero-torsion condition.

DFTT-01/94
 hep-th/9402033
 February 1994

e-mail addresses: decnet=31890::castellani;
 internet= castellani@to.infn.it

We describe in this Letter a geometric procedure to gauge the quantum Poincaré algebra found in ref. [1]. The lagrangian we obtain is a generalization of the Einstein-Cartan lagrangian, and has the same kind of symmetries (now q -deformed symmetries) as its classical counterpart: it is invariant under local Lorentz rotations and q -diffeomorphisms.

As one could expect, the differential calculus on the q -deformed Poincaré group is the correct framework for the program of finding a q -generalization of Einstein gravity. It was not obvious that this program could be carried to the end: in fact it can be done. We refer to [1] for most of the technicalities regarding the inhomogeneous quantum groups $ISO_q(N)$ and their differential calculus. Here we concentrate directly on the $ISO_q(3, 1)$ quantum Lie algebra, and discuss its gauging. The method we follow is a natural q -extension of the geometric procedure described in [2] for classical gauge and (super)gravity theories ¹.

The starting point is the q -algebra $ISO_q(3, 1)$ of ref. [1]:

$$[\chi_{ab}, \chi_{cd}] = C_{bc}\chi_{ad} + C_{ad}\chi_{bc} - C_{bd}\chi_{ac} - C_{ac}\chi_{bd} \quad (1)$$

$$\begin{aligned} [\chi_{12}, \chi_a]_{q^{-1}} &= q^{-\frac{1}{2}}C_{2a}\chi_1 - q^{-\frac{1}{2}}C_{1a}\chi_2 \\ [\chi_{13}, \chi_a]_{q^{-1}} &= q^{-\frac{1}{2}}C_{3a}\chi_1 - q^{-\frac{1}{2}}C_{1a}\chi_3 \\ [\chi_{14}, \chi_a] &= C_{4a}\chi_1 - C_{1a}\chi_4 \\ [\chi_{23}, \chi_a] &= C_{3a}\chi_2 - C_{2a}\chi_3 \\ [\chi_{24}, \chi_a]_q &= q^{\frac{1}{2}}C_{4a}\chi_2 - q^{\frac{1}{2}}C_{2a}\chi_4 \\ [\chi_{34}, \chi_a]_q &= q^{\frac{1}{2}}C_{4a}\chi_3 - q^{\frac{1}{2}}C_{3a}\chi_4 \end{aligned} \quad (2)$$

$$\begin{aligned} [\chi_1, \chi_2]_{q^{-1}} &= 0, & [\chi_1, \chi_3]_{q^{-1}} &= 0 \\ [\chi_1, \chi_4]_{q^{-2}} &= 0, & [\chi_2, \chi_3] &= 0 \\ [\chi_2, \chi_4]_{q^{-1}} &= 0, & [\chi_3, \chi_4]_{q^{-1}} &= 0 \end{aligned} \quad (3)$$

where $[A, B]_s \equiv AB - sBA$. The subalgebra spanned by the Lorentz generators χ_{ab} ($= -\chi_{ba}$) is *classical*; the deformation parameter q appears only in the commutation relations (2) and (3), involving the momenta χ_a . The metric C_{ab} is given by

$$C_{ab} = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix} \quad (4)$$

with Lorentz signature $(+, +, +, -)$. Only in the classical limit $q \rightarrow 1$ can one redefine the generators so as to diagonalize (4). The fact that the metric is diagonal

¹the so-called “group manifold approach” was initiated in ref.s [3].

in the indices 2,3 (and not completely antidiagonal as for the q -groups defined in [4]) is due to the existence of a particular real form on $SO_q(4; \mathbf{C})$. This real form, first discussed in ref. [5] for the uniparametric q -groups $SO_q(2n; \mathbf{C})$, was extended to the multiparametric case and to $ISO_q(2n, \mathbf{C})$ in [1], and allows to redefine antihermitian (linear combinations of the) generators, bringing the antidiagonal metric of ref. [4] in the hybrid form (4).

The algebra (1)-(3) was obtained in ref. [1] via a consistent projection from the q -Lie algebra of a particular multiparametric deformation of $SO(6)$, for which the R matrix takes a very simple form: it is diagonal and satisfies $\hat{R}^2 = 1$, with $\hat{R} \equiv PR$ ($\hat{R}^{ab}_{cd} \equiv R^{ba}_{cd}$).

The q -Lie algebra (1)-(3) has the form

$$\chi_i \chi_j - \Lambda^{kl}_{ij} \chi_k \chi_l = \mathbf{C}_{ij}{}^k \chi_k \quad (5)$$

where $i, j, \dots$ are adjoint indices running on the 10 values corresponding to the indices (a, ab) of the generators of $ISO_q(3, 1)$. The non-vanishing components of the braiding matrix Λ and the structure constants $\mathbf{C}$, implicitly defined by (1)-(3), are given below (no sum on repeated indices):

$$\Lambda^{ab}{}^{cd}_{ef}{}_{gh} = \delta_g^a \delta_h^b \delta_e^c \delta_f^d \quad (6)$$

$$\Lambda^{a}{}^{bc}{}_{de}{}^f = (\alpha_{de})^2 \delta_f^a \delta_d^b \delta_e^c \quad (7)$$

$$\Lambda^{bc}{}^a{}_f{}_{de} = (\alpha_{de})^{-2} \delta_f^a \delta_d^b \delta_e^c \quad (8)$$

$$\Lambda^{ab}{}_{cd} = \beta_{cd} \delta_d^a \delta_c^b \quad (9)$$

$$\mathbf{C}_{ab}{}^{cd}{}_{ef} = \frac{1}{4} [C_{ad} \delta_b^e \delta_c^f + C_{bc} \delta_a^e \delta_d^f - C_{ac} \delta_b^e \delta_d^f - C_{bd} \delta_a^e \delta_c^f] - (e \leftrightarrow f) \quad (10)$$

$$\mathbf{C}_{ab}{}^c{}_d = \frac{1}{2} \alpha_{ab} (C_{bc} \delta_a^d - C_{ac} \delta_b^d) \quad (11)$$

$$\mathbf{C}_c{}^{ab}{}_d = -\frac{1}{2} \alpha_{ab}^{-1} (C_{bc} \delta_a^d - C_{ac} \delta_b^d) \quad (12)$$

with

$$\alpha_{12} = \alpha_{13} = q^{-\frac{1}{2}}, \quad \alpha_{24} = \alpha_{34} = q^{\frac{1}{2}}, \quad \alpha_{14} = \alpha_{23} = 1 \quad (13)$$

$$\beta_{12} = \beta_{13} = \beta_{24} = \beta_{34} = q^{-1}, \quad \beta_{14} = q^{-2}, \quad \beta_{23} = 1, \quad \beta_{ab} = \beta_{ba}^{-1} \quad (14)$$

Note that the Λ tensor has unit square, i.e.

$$\Lambda^{ij}{}_{kl} \Lambda^{kl}{}_{mn} = \delta_m^i \delta_n^j \quad (15)$$

so that the algebra in (1)-(3) is a *minimal* deformation of $ISO(3, 1)$. Deformations of Lie algebras whose braiding matrix has unit square were considered some time ago by Gurevich [6].

The Λ and $\mathbf{C}$ components in (6)-(12) satisfy the following conditions ²

$$\mathbf{C}_{ri}{}^n \mathbf{C}_{nj}{}^s - \Lambda^{kl}{}_{ij} \mathbf{C}_{rk}{}^n \mathbf{C}_{nl}{}^s = \mathbf{C}_{ij}{}^k \mathbf{C}_{rk}{}^s \quad (q\text{-Jacobi identities}) \quad (16)$$

$$\Lambda^{nm}{}_{ij} \Lambda^{ik}{}_{rp} \Lambda^{js}{}_{kq} = \Lambda^{nk}{}_{ri} \Lambda^{ms}{}_{kj} \Lambda^{ij}{}_{pq} \quad (\text{Yang-Baxter}) \quad (17)$$

$$\mathbf{C}_{mn}{}^i \Lambda^{ml}{}_{rj} \Lambda^{ns}{}_{lk} + \Lambda^{il}{}_{rj} \mathbf{C}_{lk}{}^s = \Lambda^{pq}{}_{jk} \Lambda^{is}{}_{lq} \mathbf{C}_{rp}{}^l + \mathbf{C}_{jk}{}^m \Lambda^{is}{}_{rm} \quad (18)$$

$$\mathbf{C}_{rk}{}^m \Lambda^{ns}{}_{ml} = \Lambda^{ij}{}_{kl} \Lambda^{nm}{}_{ri} \mathbf{C}_{mj}{}^s \quad (19)$$

These are the “bicovariance conditions”, see ref.s [7, 8, 9], necessary for the existence of a bicovariant differential calculus (see also the discussion in Appendix B of [1]). Whenever we have a set of matrices $\Lambda^{ij}{}_{kl}$, $\mathbf{C}_{ij}{}^k$ satisfying (16)-(19) we can construct a differential calculus on the quantum group $Fun_q(M_i{}^j)$, generated by the elements (adjoint representation of the q -groups) $M_i{}^j$ satisfying the “ ΛMM ” relations:

$$M_i{}^j M_r{}^q \Lambda^{ir}{}_{pk} = \Lambda^{jq}{}_{ri} M_p{}^r M_k{}^i \quad (20)$$

Consistency of these relations is ensured by the QYB equations (17). One can define in the usual way a coproduct $\Delta(M_i{}^j) = M_i{}^k \otimes M_k{}^j$ and a counit $\varepsilon(M_i{}^j) = \delta_i^j$. When $\Lambda^2 = 1$ one can also define a coinverse $\kappa(M_i{}^j)$ with $\kappa^2 = 1$ (This is done by enlarging the algebra $Fun_q(M_i{}^j)$, see Appendix B of [1]).

The generators χ_i of the q -Lie algebra (5) are functionals on $Fun_q(M_i{}^j)$:

$$\chi_j(M_i{}^k) = \mathbf{C}_{ij}{}^k \quad (21)$$

We recall that products of functionals are defined via the coproduct Δ , i.e. $\chi_i \chi_j \equiv (\chi_i \otimes \chi_j) \Delta$, whereas functionals act on products as $\chi_i(ab) = \Delta'(\chi_i)(a \otimes b)$, $a, b \in Fun_q(M_i{}^j)$ (see below the definition of Δ').

Next we introduce new functionals $f^i{}_l$ via their action on the basis $M_k{}^j$:

$$f^i{}_l(M_k{}^j) = \Lambda^{ij}{}_{kl} \quad (22)$$

The co-structures of χ and f are given by:

$$\Delta'(\chi_i) = \chi_j \otimes f^j{}_i + I' \otimes \chi_i \quad (23)$$

$$\varepsilon'(\chi_i) = 0 \quad (24)$$

$$\kappa'(\chi_i) = -\chi_j \kappa'(f^j{}_i) \quad (25)$$

$$\Delta'(f^i{}_j) = f^i{}_k \otimes f^k{}_j \quad (26)$$

$$\varepsilon'(f^i{}_j) = \delta_j^i \quad (27)$$

$$\kappa'(f^i{}_j) = f^i{}_j \circ \kappa \quad (28)$$

The algebra generated by the χ and f is a Hopf algebra (the χ, f and f, f commutations are given in [8, 9, 1]), and defines a bicovariant differential calculus

²this can be checked directly without too much effort. In [1] a general proof is given for all $ISO_q(N)$ obtained by the projective method.

on the q -group generated by the M_i^j elements. For example, one can introduce left-invariant one-forms ω^i as duals to the “tangent vectors” χ_i , an exterior product

$$\omega^i \wedge \omega^j \equiv \omega^i \otimes \omega^j - \Lambda^{ij}{}_{kl} \omega^k \otimes \omega^l, \quad (29)$$

an exterior derivative on $Fun_q(M_i^j)$ as

$$da = (id \otimes \chi_i) \Delta(a) \omega^i, \quad a \in Fun_q(M_i^j) \quad (30)$$

and so on. The commutations between one-forms and elements $a \in Fun_q(M_i^j)$ are given by:

$$\omega^i a = (id \otimes f^i{}_j) \Delta(a) \quad (31)$$

The exterior derivative can be extended to the (left-invariant) one-forms via the deformed Cartan-Maurer equations [7, 9]

$$d\omega^i + C_{jk}{}^i \omega^j \wedge \omega^k = 0 \quad (32)$$

The C structure constants appearing in the Cartan-Maurer equations are related to the $\mathbf{C}$ constants of the q -Lie algebra as [9]:

$$\mathbf{C}_{jk}{}^i = C_{jk}{}^i - \Lambda^{rs}{}_{jk} C_{rs}{}^i \quad (33)$$

In the particular case $\Lambda^2 = I$ it is not difficult to see that $C = \frac{1}{2}\mathbf{C}$.

The procedure we have advocated in refs [10] for the “gauging” of quantum groups essentially retraces the steps of the group-geometric method for the gauging of usual Lie groups, described for instance in refs [2].

We consider one-forms ω^i which are not left-invariant any more, so that the Cartan-Maurer equations are replaced by:

$$R^i = d\omega^i + C_{jk}{}^i \omega^j \wedge \omega^k \quad (34)$$

where the curvatures R^i are now non-vanishing, and satisfy the q -Bianchi identities:

$$dR^i - C_{jk}{}^i R^j \wedge \omega^k + C_{jk}{}^i \omega^j \wedge R^k = 0 \quad (35)$$

due to the Jacobi identities on the structure constants $C_{ij}{}^k$ [9]. As in the classical case we can write the q -Bianchi identities as $\nabla R^i = 0$ (these define the covariant derivative ∇).

Consider now the definition (34) of the curvature R^i , and apply it to the q -Poincaré algebra of (1)-(3): the one-forms dual to χ_{ab} , χ_a are respectively denoted by ω^{ab} , V^a and the corresponding curvatures read (we omit wedge symbols):

$$R^{ab} = d\omega^{ab} + C_{cd}{}^{ab} \omega^{cd} \quad (36)$$

$$R^a = dV^a + \alpha_{af} C_{fb}{}^a \omega^f V^b \quad (37)$$

where $V_a \equiv C_{ab}V^b$, α_{af} and C_{ab} are given in (13) and (4), and we used $C_{ij}{}^k = \frac{1}{2}\mathbf{C}_{ij}{}^k$. We have also rescaled ω^{ab} by a factor $\frac{1}{2}$ to obtain standard normalizations. R^{ab} is the q -Lorentz curvature, coinciding with the classical one (as a function of ω^{ab}), and R^a is the q -deformed torsion.

The Bianchi identities, deduced from (35), are:

$$dR^{ab} - C_{fe}R^{af}\omega^{eb} + C_{fe}\omega^{af}R^{eb} = 0 \quad (38)$$

$$dR^a + \alpha_{af}C_{fb}R^{af}V^b - \alpha_{af}C_{fb}\omega^{af}R^b = 0 \quad (39)$$

From the definition (29) of the exterior product we see that for $\Lambda^2 = I$ the one-forms ω^i q -commute as:

$$\omega^i\omega^j = -\Lambda^{ij}{}_{kl}\omega^k\omega^l \quad (40)$$

Inserting the Λ tensor corresponding to (6)-(9) we find:

$$\begin{aligned} V^a\omega^{12} &= -q^{-1}\omega^{12}V^a \\ V^a\omega^{13} &= -q^{-1}\omega^{13}V^a \\ V^a\omega^{14} &= -\omega^{14}V^a \\ V^a\omega^{23} &= -\omega^{23}V^a \\ V^a\omega^{24} &= -q\omega^{24}V^a \\ V^a\omega^{34} &= -q\omega^{34}V^a \end{aligned} \quad (41)$$

$$\begin{aligned} V^2V^1 &= -q^{-1}V^1V^2 \\ V^3V^1 &= -q^{-1}V^1V^3 \\ V^4V^1 &= -q^{-2}V^1V^4 \\ V^3V^2 &= -V^2V^3 \\ V^4V^2 &= -q^{-1}V^2V^4 \\ V^4V^3 &= -q^{-1}V^3V^4 \end{aligned} \quad (42)$$

and usual anticommutations between the ω^{ab} (components of the Lorentz spin connection). The exterior product of two identical one-forms vanishes (this is not true in general when $\Lambda^2 \neq I$). As a consequence, the exterior product of five vielbeins is zero.

We are now ready to write the lagrangian for the q -gravity theory based on $ISO_q(3,1)$. The lagrangian looks identical to the classical one, i.e.:

$$\mathcal{L} = R^{ab}V^cV^d\varepsilon_{abcd} \quad (43)$$

The Lorentz curvature R^{ab} , although defined as in the classical case, has non-trivial commutations with the q -vielbein:

$$V^aR^{12} = q^{-1}R^{12}V^a$$

$$\begin{aligned}
V^a R^{13} &= q^{-1} R^{13} V^a \\
V^a R^{14} &= R^{14} V^a \\
V^a R^{23} &= R^{23} V^a \\
V^a R^{24} &= q R^{24} V^a \\
V^a R^{34} &= q R^{34} V^a
\end{aligned} \tag{44}$$

deducible from the definition (36). As in ref. [10, 9], we make the assumption that the commutations of $d\omega^i$ with the one-forms ω^l are the same as those of $C_{jk}^i \omega^j \omega^k$ with ω^l , i.e. the same as those valid for $R^i = 0$. For the definition of ε_{abcd} in (43) see below.

We discuss now the notion of q -diffeomorphisms. It is known that there is a consistent q -generalization of the Lie derivative (see ref.s [9, 11, 12, 1]) which can be expressed as in the classical case as:

$$\ell_V = i_V d + di_V \tag{45}$$

where i_V is the q -contraction operator defined in ref.s [9, 11], with the following properties:

$$\begin{aligned}
i) \quad & i_V(a) = 0, \quad a \in A, \quad V \text{ generic tangent vector} \\
ii) \quad & i_{t_i} \omega^j = \delta_i^j I \\
iii) \quad & i_{t_i}(\theta \wedge \omega^k) = i_{t_r}(\theta) \omega^l \Lambda^{rk}_{li} + (-1)^p \theta \delta_i^k, \quad \theta \text{ generic } p\text{-form} \\
iv) \quad & i_V(a\theta + \theta') = ai_V(\theta) + i_V\theta', \quad \theta, \theta' \text{ generic forms} \\
v) \quad & i_{\lambda V} = \lambda i_V, \quad \lambda \in \mathbf{C} \\
vi) \quad & i_{\varepsilon V}(\theta) = i_V(\theta)\varepsilon, \quad \varepsilon \in A
\end{aligned} \tag{46}$$

As a consequence, the q -Lie derivative satisfies:

$$\begin{aligned}
i) \quad & \ell_V a = i_V(da) \equiv V(a) \\
ii) \quad & \ell_V d = d\ell_V \\
iii) \quad & \ell_V(\lambda\theta + \theta') = \lambda\ell_V(\theta) + \ell_V(\theta') \\
iv) \quad & \ell_{\varepsilon V}(\theta) = (\ell_V\theta)\varepsilon - (-1)^p i_V(\theta)d\varepsilon, \quad \theta \text{ generic } p\text{-form} \\
v) \quad & \ell_{t_i}(\theta \wedge \omega^k) = (\ell_{t_r}\theta) \wedge \omega^l \Lambda^{rk}_{li} + \theta \wedge \ell_{t_i}\omega^k
\end{aligned} \tag{47}$$

In analogy with the classical case, we define the q -diffeomorphism variation of the fundamental field ω^i as

$$\delta\omega^k \equiv \ell_{\varepsilon^{i_{t_i}}}\omega^k \tag{48}$$

where according to iv) in (47):

$$\ell_{\varepsilon^{i_{t_i}}}\omega^k = (i_{t_i}d\omega^k + di_{t_i}\omega^k)\varepsilon^i + d\varepsilon^k = (i_{t_i}d\omega^k)\varepsilon^i + d\varepsilon^k \tag{49}$$

As in the classical case, there is a suggestive way to write this variation:

$$\ell_{\varepsilon^{i_{t_i}}}\omega^k = i_{\varepsilon^{i_{t_i}}}R^k + \nabla\varepsilon^k \tag{50}$$

where

$$\begin{aligned}\nabla \varepsilon^k &\equiv d\varepsilon^k - C_{rs}{}^k i_{t_i}(\omega^r \wedge \omega^s) \varepsilon^i = \\ &d\varepsilon^k - C_{rs}{}^k \varepsilon^r \omega^s + C_{rs}{}^k \omega^r \varepsilon^s\end{aligned}\quad (51)$$

Proof: use the Bianchi identities (35) and iii) in (46).

Notice that if we postulate:

$$\begin{aligned}\Lambda^{rk}{}_{li} \omega^l \varepsilon^i &= \varepsilon^r \omega^k \\ \Lambda^{rk}{}_{li} \omega^l \wedge d\varepsilon^i &= -d\varepsilon^r \wedge \omega^k\end{aligned}\quad (52)$$

we find

$$\delta(\omega^j \wedge \omega^k) = \delta\omega^j \wedge \omega^k + \omega^j \wedge \delta\omega^k \quad (53)$$

i.e. a rule that any “sensible” variation law should satisfy. To prove (53) use iv) and v) of (47). The q -commutations (52) were already proposed in [10] in the context of q -gauge theories. A consequence of (52) are the following commutations between the variation parameter and $d\omega^i$:

$$\Lambda^{rk}{}_{li} d\omega^l \varepsilon^i = \varepsilon^r d\omega^k \quad (54)$$

As discussed in ref.s [10], it is consistent to postulate that R^i has the same commutations with ε^j as $d\omega^i$:

$$\Lambda^{rk}{}_{li} R^l \varepsilon^i = \varepsilon^r R^k \quad (55)$$

We have now all the tools we need to investigate the invariances of the q -gravity lagrangian (43). The result will be analogous to the classical one: after imposing the horizontality conditions

$$i_{t_{ab}} R^{cd} = i_{t_{ab}} R^c = 0 \quad (56)$$

along the Lorentz directions one finds that, *provided* the ε tensor in (43) is appropriately defined, the lagrangian is invariant under q -diffeomorphisms and local Lorentz rotations. Note that, as in the $q = 1$ case, the horizontality conditions (56) can be obtained as field equations (see later).

We first consider Lorentz rotations. Under these, the curvature R^{ab} and the vielbein V^c transform as:

$$\delta R^{ab} \equiv \ell_{\varepsilon^{gh} t_{gh}} R^{ab} = \mathbf{C}_{ef}{}^{ab}{}_{gh} R^{ef} \varepsilon^{gh} - \mathbf{C}_{gh}{}^{ab}{}_{ef} \varepsilon^{gh} R^{ef} \quad (57)$$

$$\delta V^c \equiv \ell_{\varepsilon^{gh} t_{gh}} V^c = -\mathbf{C}_{ef}{}^c{}_{g} \varepsilon^{ef} V^g + \mathbf{C}_g{}_{ef}{}^c V^g \varepsilon^{ef} \quad (58)$$

To obtain these variations, use the definition (45), iv) of (47), the Bianchi identity (38) and the horizontality conditions (56).

Now we have the lemma:

$$\delta\mathcal{L} = [(\delta R^{ab})V^cV^d + R^{ab}(\delta V^c)V^d + R^{ab}V^c(\delta V^d)]\varepsilon_{abcd} \quad (59)$$

Proof: use v) of (47) and the first of (52).

Inserting the variations (57) and (58) inside (59) we find, after ordering the terms as εRVV with (52) and (55):

$$\delta\mathcal{L} = 2(\mathbf{C}_{ef\ gh}^{ab}\varepsilon_{abcd} - \mathbf{C}_{gh\ c}^s\varepsilon_{efsd} - \mathbf{C}_{rs\ d}^p\Lambda^q{}_{gh\ c}{}^{rs}\varepsilon_{efqp})\varepsilon^{gh}R^{ef}V^cV^d \quad (60)$$

Using the explicit form of the Λ and $\mathbf{C}$ tensors into (6)-(12) and imposing $\delta\mathcal{L} = 0$, we find a set of equations for the ε_{abcd} tensor. These can in fact be solved and yield:

$$\begin{aligned} \varepsilon_{1234} &= 1, & \varepsilon_{1243} &= -q, & \varepsilon_{1324} &= -1, & \varepsilon_{1342} &= q, \\ \varepsilon_{1423} &= q^{\frac{3}{2}}, & \varepsilon_{1432} &= -q^{\frac{3}{2}}, & \varepsilon_{2134} &= -1, & \varepsilon_{2143} &= q, \\ \varepsilon_{3124} &= 1, & \varepsilon_{3142} &= -q, & \varepsilon_{4123} &= -q^{\frac{3}{2}}, & \varepsilon_{4132} &= q^{\frac{3}{2}}, \\ \varepsilon_{2314} &= q^{\frac{1}{2}}, & \varepsilon_{2341} &= -q^{\frac{5}{2}}, & \varepsilon_{2413} &= -q^2, & \varepsilon_{2431} &= q^3, \\ \varepsilon_{3214} &= -q^{\frac{1}{2}}, & \varepsilon_{3241} &= q^{\frac{5}{2}}, & \varepsilon_{4213} &= q^2, & \varepsilon_{4231} &= -q^3, \\ \varepsilon_{3412} &= q^2, & \varepsilon_{3421} &= -q^3, & \varepsilon_{4312} &= -q^2, & \varepsilon_{4321} &= q^3 \end{aligned} \quad (61)$$

Consider next the variation of $\mathcal{L}$ under q -diffeomorphisms, i.e.:

$$\delta\mathcal{L} = \ell_{\varepsilon^g t_g}\mathcal{L} = (\ell_{t_g}\mathcal{L})\varepsilon^g - (i_{t_g}\mathcal{L})d\varepsilon^g = \quad (62)$$

$$= d[i_{t_g}(R^{ab}V^cV^d\varepsilon_{abcd})\varepsilon^g] + i_{t_g}[d(R^{ab}V^cV^d\varepsilon_{abcd})]\varepsilon^g \quad (63)$$

Then the variation $\delta\mathcal{L}$ is a total derivative if

$$d(R^{ab}V^cV^d\varepsilon_{abcd}) = 0 \quad (64)$$

After using the expression for dR^{ab} given by the Bianchi identity (38) and the torsion definition (37) to find dV^a (note that $R^{ab}R^cV^d\varepsilon_{abcd} = 0$ because of horizontality of R^{ab} , R^c and the vanishing of the product of five vielbeins), eq. (64) yields a set of conditions on ε_{abcd} . These conditions in fact *coincide* with those found to ensure local Lorentz invariance of the q -lagrangian. This is not a miracle: indeed we could have computed the Lorentz variation of $\mathcal{L}$ in the same way as in (63); the total derivative term would have vanished because $i_{t_{gh}}(R^{ab}V^cV^d) = 0$ (horizontality of R^{ab}), and we would have found again eq. (64) as a condition for $\delta\mathcal{L} = 0$.

Thus the lagrangian (43) with the ε_{abcd} tensor as given in (61) is invariant (up to a total derivative) also under q -diffeomorphisms.

We discuss now the algebra of q -Lie derivatives. We have the theorem, analogous to the classical one:

$$\ell_{t_i}\ell_{t_j} - \Lambda^{kl}{}_{ij}\ell_{t_k}\ell_{t_l} = \ell_{(\mathbf{C}_{ij}{}^n{}_{-R^n{}_{ij})t_n}} \quad (65)$$

with

$$R^i \equiv R^i_{jk} \omega^j \wedge \omega^k \quad (66)$$

$$\mathbf{R}^i_{jk} \equiv R^i_{jk} - \Lambda^{rs}_{jk} R^i_{rs} \quad (67)$$

As for the structure constants, we have $R^i_{jk} = \frac{1}{2} \mathbf{R}^i_{jk}$ when $\Lambda^2 = 1$. The proof of the composition law (65) is computational: one applies its left-hand side to ω^k , and uses the properties of the q -Lie derivative. *Hint 1*: rewrite the Lie derivative of ω^k as:

$$\ell_{t_j} \omega^k = (\mathbf{C}_{rj}{}^k - \mathbf{R}^k_{rj}) \omega^r \quad (68)$$

Hint 2: use the following expression for Λ^{ij}_{kl} (no sums on repeated indices) :

$$\Lambda^{ij}_{kl} = [kl] \delta^i_l \delta^j_k \quad (69)$$

and the identities

$$[kl] = \frac{1}{[lk]} \quad (70)$$

$$\mathbf{C}_{rk}{}^s [lk] [lr] = \mathbf{C}_{rk}{}^s [ls] \quad (71)$$

due to $\Lambda^2 = 1$ and the bicovariance condition (19).

From the q -algebra (65) it is not difficult to find the following composition law for q -variations:

$$\ell_{\varepsilon_2^i t_i} \ell_{\varepsilon_1^j t_j} - (1 \leftrightarrow 2) = \ell_{(\mathbf{C}_{ij}{}^n - \mathbf{R}^n_{ij}) \varepsilon_2^i \varepsilon_1^j t_n} \quad (72)$$

The order of the factors is important in the composite parameter $(\mathbf{C}_{ij}{}^n - \mathbf{R}^n_{ij}) \varepsilon_2^i \varepsilon_1^j$. Note also that

$$(\mathbf{C}_{ij}{}^n - \mathbf{R}^n_{ij}) \varepsilon_2^i \varepsilon_1^j = \frac{1}{2} (\mathbf{C}_{ij}{}^n - \mathbf{R}^n_{ij}) (\varepsilon_2^i \varepsilon_1^j - \varepsilon_1^i \varepsilon_2^j) \quad (73)$$

if we postulate the commutation rule

$$\varepsilon_1^i \varepsilon_2^j = \Lambda^{ij}_{kl} \varepsilon_2^k \varepsilon_1^l \quad (74)$$

Indeed $(\mathbf{C}_{ij}{}^n - \mathbf{R}^n_{ij}) \Lambda^{ij}_{kl} = -(\mathbf{C}_{kl}{}^n - \mathbf{R}^n_{kl})$ (due to $\Lambda^2 = 1$). Then the composite parameter is explicitly $(1 \leftrightarrow 2)$ antisymmetric.

Let us derive the equations of motion corresponding to the q -lagrangian (43). We assume the same variational rule as with the Lie derivative. The q -Einstein equations are obtained by varying the vielbein in $\mathcal{L}$:

$$\delta \mathcal{L} = R^{ab} (\delta V^c V^d + V^c \delta V^d) \varepsilon_{abcd} = 0 \quad (75)$$

Postulating that $\delta \omega^i$ has the same commutations as ω^i , and noticing that

$$\varepsilon_{abef} = -\Lambda^{cd}_{ef} \varepsilon_{abcd} \quad (76)$$

(use the explicit entries in (9) and (61), or notice that since ε_{abcd} multiplies $V^c V^d$ in (43), it must be Λ -antisymmetric in the indices c,d), one arrives at:

$$R^{ab} V^e \varepsilon_{abef} = 0 \quad (77)$$

The q -Einstein equations are found as in the classical case: expand (77) along three vielbeins:

$$R^{ab}{}_{cd} V^c V^d V^e \varepsilon_{abef} = 0, \quad (78)$$

multiply by another vielbein V^g and use:

$$\varepsilon^{cdeg} V^1 V^2 V^3 V^4 \equiv V^c V^d V^e V^g \quad (79)$$

(N.B.: the entries of ε^{abcd} are different from those of ε_{abcd}) so that finally we have:

$$R^{ab}{}_{cd} \varepsilon^{cdeg} \varepsilon_{abef} = 0 \quad (80)$$

The contraction of the two alternating tensors yields a q -weighted product of Kronecker deltas. We leave to the reader to find the final form of the q -Einstein equations. Expanding (77) on ωVV , $V\omega V$ and $\omega\omega V$ yields instead the horizontality condition on R^{ab} .

The torsion equation is obtained by varying (43) in the spin connection ω^{ab} . The final result is again an equation that formally looks identical to the classical one:

$$R^c V^d \varepsilon_{abcd} = 0 \quad (81)$$

As for $q = 1$, this equation implies that the torsion vanishes as a two-form: $R^c = 0$ (hence also horizontality of R^c).

Note 1: The q -volume 4-form $V^1 V^2 V^3 V^4 = f(q) V^a V^b V^c V^d \varepsilon_{abcd}$ (with $[f(q)]^{-1} = \varepsilon^{abcd} \varepsilon_{abcd} = 2q^{-\frac{3}{2}}(2q^{-\frac{3}{2}} + q^{-1} + 2q^{-\frac{1}{2}} + 2 + 2q^{\frac{1}{2}} + q + 2q^{\frac{3}{2}})$) is invariant under Lorentz rotations and, up to a total derivative, under q -diffeomorphisms. The proof is similar to the one used for the lagrangian. This means that the q -symmetries allow a cosmological constant term $V^a V^b V^c V^d \varepsilon_{abcd}$.

Note 2: It would be worthwhile to give a recipe for extracting numbers out of a q -theory of the kind discussed in this Letter. A possible way of doing this would be to find a consistent definition of path-integral on q -commuting fields, leading to **C**-number amplitudes. On this question, see for example [13].

Acknowledgements

It is a pleasure to thank the Centro Brasileiro de Pesquisas Físicas of Rio de Janeiro, where this work was completed, for its warm hospitality.

References

- [1] L. Castellani, *Differential calculus on $ISO_q(N)$, quantum Poincaré algebra and q -gravity*, Torino preprint DFTT-70/93, hep-th 9312179.
- [2] L. Castellani, R. D'Auria and P. Fré, *Supergravity and Superstrings: a geometric perspective* (World Scientific, Singapore, 1991); L. Castellani, Int. J. Mod. Phys. **A7** (1992) 1583.
- [3] Y. Ne'eman and T. Regge, Riv. Nuovo Cim. **5** (1978) 1; Phys.Lett.**74B** (1978) 54; A. D'Adda, R. D'Auria, P. Fré, T. Regge, Riv. Nuovo Cim. **6** (1980) 3.
- [4] L.D. Faddeev, N.Yu. Reshetikhin and L.A. Takhtajan, Algebra and Analysis, **1** (1987) 178.
- [5] E. Celeghini, R. Giachetti, A. Reyman, E. Sorace, M. Tarlini, $SO_q(n+1, n-1)$ as a real form of $SO_q(2n, \mathbf{C})$, Lett. Math. Phys. **23** (1991) 45.
- [6] D.I. Gurevich, Soviet Math. Dokl., **33** (1986) 758.
- [7] S.L. Woronowicz, Publ. RIMS, Kyoto Univ., Vol. **23**, 1 (1987) 117; Commun. Math. Phys. **111** (1987) 613 and Commun. Math. Phys. **122**, (1989) 125.
- [8] D. Bernard, *Quantum Lie algebras and differential calculus on quantum groups*, Progress of Theor. Phys. Suppl. **102** (1990) 49; Phys. Lett. **260B** (1991) 389.
- [9] P. Aschieri and L. Castellani, Int. Jou. Mod. Phys. **A8** (1993) 1667.
- [10] L. Castellani, Phys. Lett. **B292** (1992) 93; $U_q(N)$ gauge theories, Torino preprint DFTT 74/92.
- [11] P. Aschieri, *The space of vector fields on quantum groups*, UCLA preprint UCLA/93/TEP/25.
- [12] P. Schupp, P. Watts and B. Zumino, Lett. Math. Phys. **25** (1992) 139; *Cartan calculus for Hopf algebras and quantum groups*, NSF-ITP-93-75, LBL-34215 and UCB-PTH-93/20; P. Schupp, *Quantum groups, non-commutative differential geometry and applications*, Ph.D. Thesis, LBL-34942 and UCB-PTH-93/35.
- [13] L. Baulieu and E.G. Floratos, Phys. Lett. **B258** (1991) 171.